

First record and range extension of the freshwater limpet *Gundlachia radiata* (Guilding, 1828) (Mollusca: Gastropoda: Planorbidae) from southeast Brazil

Luiz Eduardo Macedo de Lacerda, Igor Christo Miyahira and Sonia Barbosa dos Santos*

Universidade do Estado do Rio de Janeiro, Instituto de Biologia Roberto Alcântara Gomes, Programa de Pós-graduação em Ecologia e Evolução, Departamento de Zoologia, Laboratório de Malacologia Limnica e Terrestre. Rua São Francisco Xavier, 524, sala 525/2. CEP 20550-900. Rio de Janeiro, RJ, Brazil.

* Corresponding author. E-mail: sbsantos@uerj.br

ABSTRACT: We report the first records of *Gundlachia radiata* (Guilding, 1828), a freshwater limpet, from the state of Rio de Janeiro, which extends its distribution to Southeast Brazil.

Gundlachia Pfeiffer, 1849, is one of the seven genera of the traditional Ancyliidae (Hubendick 1967) that occur in the Neotropical region (Santos 2003), now included in the Ancylini tribe of Planorbidae (Bouchet & Rocroi 2005; Albrecht *et al.* 2007). This genus contains the greatest number of species among “Neotropical ancyliids” (Lanzer 1996; Santos 2003; Ovando *et al.* 2011). The type locality of *Gundlachia radiata* (Guilding, 1828) is Saint Vincent, an island in the Caribbean Sea (Central America). The species is known for 24 countries in the Americas (Ovando *et al.* 2011). In Brazil, it is found in the North region to the states of Amazonas (Irmiler 1975) and Pará (Lanzer 1996), in the Northern region to the state of Paraíba (Col. Mol. UERJ 54), Pernambuco (Lanzer 1996), Alagoas (Col. Mol. UERJ 25), and to the Central-Western region to the state of Goiás (Thiengo *et al.* 2005).

We report the first records of *Gundlachia radiata* (Figure 1) for the state of Rio de Janeiro (SRJ), and extend the species’ geographic distribution to Southeast Brazil (Figure 2). We also present data on morphological variations in shell and soft-parts.

The material studied was collected from the municipalities of Rio de Janeiro and Nova Iguaçu in Rio de Janeiro state, Brazil (Figure 2). Fieldwork was conducted

under legal authorization (SISBIO 10812-1 and 23607-2). Species identification was done under stereomicroscope, and was based on shell shape and size, teleoconch and protoconch microsculpture, mantle pigmentation pattern and adductor muscle scars. Electron scanning microscopy images were also taken. Illustrations were done under *camera lucida*. Shell measurements and diagnostic characters were taken according Hubendick (1964), Lanzer (1996) and Santos (2003). Samples are housed in the Malacological Collection of the Universidade do Estado do Rio de Janeiro (Col. Mol. UERJ).

Examined material: Brazil, *Rio de Janeiro*: Rio de Janeiro (Santa Cruz, São Francisco channel, 22°51’43” S, 43°40’48” W), 10 animals (shells and soft parts), 13.VIII.2010, (Col. Mol. UERJ 8363). Nova Iguaçu (Guandu river, 22°48’22” S, 43°37’33” W), 49 animals (shells and soft parts), 26.IV.2011, (Col. Mol. UERJ 8379).

Although the collecting sites have different names (São Francisco Channel and Guandu River), both are in the same hydrographic catchment. Guandu River site is 11 km upriver from the São Francisco Channel site. The collection site in São Francisco channel has sandy bottoms and intense water flow. Both margins of the river have agricultural crops and sand is extracted commercially from the river. The second collecting site, in Guandu River (Nova Iguaçu), is at a dam adjacent to a water treatment plant with low water flow. In both sites, ancyliids are found attached to *Eichornia* sp. (water hyacinth).

The 59 specimens show an elliptical shell opening (Figure 3A). The apex is rounded and right-posteriorly pointed; there is a shallow apical depression and up to two-thirds of the protoconch has irregular shallow pits (Figure 3). The anterior right adductor muscle scar is more elliptical than the left, with a slight curvature in the inner portion (Figure 4). All of these characters fit well with descriptions of *Gundlachia radiata* (Hubendick 1967; Lanzer 1996; Santos 2003). All specimens from Santa Cruz, Municipality of Rio de Janeiro, have dark pigmentation concentrated on the anterior region of the mantle roof whereas the specimens from Municipality of Nova Iguaçu




FIGURE 1. Dorsal view of *Gundlachia radiata* from the municipality of Nova Iguaçu, state of Rio de Janeiro, Brazil. Scale = 1mm. Photo by L.E.M. Lacerda.




FIGURE 2. Map showing the localities in Rio de Janeiro state where *Gundlachia radiata* has been recorded. (1) Municipality of Rio de Janeiro (São Francisco Channel) and (2) municipality of Nova Iguaçu (Guandu River). Satellite image from Google Earth.

show three pigmentation patterns: 1) moderate, dark blotches on the mantle edge (55% of specimens); 2) weak, two blotches, anterior and posterior (29% of specimens); and 3) heavy, pigmentation of all of the mantle roof (16% of specimens) (Figure 4A-D). The variation in pigmentation patterns is not a good systematic character, although absence of pigmentation could be diagnostic for other genera such as *Ferrissia* Walker, 1903 and *Burnupia* Walker, 1912 (Lanzer 1996; Santos 2003).

The average measurements (mm) and standard deviation from 49 shells of *G. radiata* collected in the Municipality of Rio de Janeiro and their ranges (maximum-minimum) were 6.10 ± 1.81 (9.72-2.82) length, 3.65 ± 1.05 (5.93-1.93) width and 1.32 ± 0.40 (2.06-0.49) height; and in the Municipality of Nova Iguaçu, 4.32 ± 1.24 (6.15-2.25) length, 2.85 ± 1.00 (4.50-1.40) width and 1.16 ± 0.37 (1.80-0.60) height. Considering all specimens together, they

were 5.78 ± 1.84 (9.72-2.25) length, 3.51 ± 1.07 (5.93-1.40) width and 1.29 ± 0.39 (2.06-0.49) height.

A septum closing the aperture of the shell is commonly found in some ancylid populations, when submitted to adverse environmental conditions, preventing water loss (Aguayo 1946; Hylton-Scott 1954; Richardot 1974; Santos 2003). Such shells are said to be gundlachoid, as the genus *Gundlachia* was described based on this character (Guilding, 1828). We did not find any gundlachoid shells in the studied material although this kind of shell is common in *Gundlachia radiata* populations (Hubendick 1967; Santos 2003; Ovando *et al.* 2011). This situation was found by Ovando *et al.* (2011) in temporary ponds of Argentina, where almost 50% of specimens were gundlachoid, indicating a stressful environment for *G. radiata*; recently a septum formation was also described to *Gundlachia ticaga* Marcus & Marcus, 1962 to the state of




FIGURE 3. *Gundlachia radiata* shell. (A) Dorsal view, showing absence of radial lines on the teleoconch. Square: protoconch area. (B) Detail of the protoconch showing the irregularly arranged pits and apical depression. Photo by A. Moraes.

São Paulo, Brazil (Simone *et al.* 2012).

The present record of *Gundlachia radiata* from Rio de Janeiro and Nova Iguaçu represents the first record of the species for the state of Rio de Janeiro and a range extension to Southeast Brazil. Two hypotheses could explain this new record: it is a recent introduction, because previous surveys in the studied area (Thiengo *et al.* 2001) were

not able to find the species or, its real occurrence area has been underestimated because surveys in the state of Rio de Janeiro have focused primarily on schistosomiasis vectors (Thiengo *et al.* 2002a; b; 2004a; b; 2006). The present record increases one additional species to the list of ancylids of Rio de Janeiro state, which now reaches seven known species.




FIGURE 4. *Gundlachia radiata* from the state of Rio de Janeiro: variation on mantle pattern pigmentation (A) Specimens from Santa Cruz, Rio de Janeiro Municipality. (B-D) Specimens from Nova Iguaçu Municipality; (B) Moderate; (C) Weak and (D) Heavy. Scale = 1mm. Drawings by L.E.M. Lacerda.

ACKNOWLEDGMENTS: We thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for SBS research grant (Universal 476682/2004-5; Protax 562291/2010-5) and CNPq for LEML and ICM doctoral scholarships; Laboratório de Microscopia Eletrônica de Varredura of Universidade do Estado do Rio de Janeiro (LABMEL) for scanning microscopy images; Jéssica Beck Carneiro, Isabela Cristina Brito Gonçalves and Renata Freitas Ximenes for helping in fieldwork; Dr. Timothy Peter Moulton for the review of the final manuscript; and the anonymous reviewers for their valuable comments and suggestions to improve the quality of the paper.

LITERATURE CITED

- Aguayo, C.G. 1946. *Gundlachia* y *Hebetancylus* en Cuba. *Revista de la Sociedad Malacologica "Carlos de La Torre"* 4: 55-62.
- Albrecht, C., K. Kuhn and B. Streit. 2007. A molecular phylogeny of Planorboidea (Gastropoda, Pulmonata): insights from enhanced taxon sampling. *Zoological Scripta* 36(1): 27-39.
- Basch, P.F. 1963. A review of the recent freshwater limpet snails of North America. *Bulletin of the Museum of Comparative Zoology at Harvard College* 129: 401-461.
- Bouchet, P. and Rocroi, J.P. 2005. Classification and nomenclator of Gastropod families. *Malacologia* 47: 1-397.
- Guilding, L. 1828. Observations on the zoology of the Caribbean Islands. *Zoological Journal* 3: 527-544.
- Hubendick, B. 1967. Studies on Ancyliidae. The Australian, Pacific and Neotropical formgroups. *Göteborgs Kungl. Vetenskap-Och Vitterhets-Samhälles Handlingar Zoologie* 1: 1-52.
- Hylton-Scott, M.I. 1953. Notas sobre morfología de *Gundlachia* Pfr. (Ancyliidae). *Physis* 20: 467-473.
- Irmiler, U. 1975. Ecological studies of aquatic soil invertebrates in three inundation forests of Central-Amaonia. *Amazoniana* 5: 337-409.
- Lacerda, L.E.M., I.C. Miyahira and S.B. Santos. 2011. Shell morphology of the freshwater snail *Gundlachia ticaga* (Gastropoda, Ancyliidae) from four sites in Ilha Grande, southeastern Brazil. *Zoologia* 28(3): 334-342.
- Lacerda, L.E.M. and S.B. Santos. 2011. Mollusca, Heterobranchia, *Burnupia ingae* Lanzer, 1991 (Gastropoda, Ancyliidae): current distribution in Brazil. *Check List* 7(6): 862-864.
- Lanzer, R.M. 1996. Ancyliidae (Gastropoda: Basommatophora) na América do Sul: sistemática e distribuição. *Revista Brasileira de Zoologia* 13(1): 175-210.
- Ovando, X.M.C., L.E.M. Lacerda and S.B. Santos. 2011. Mollusca, Gastropoda, Heterobranchia, Ancyliidae, *Gundlachia radiata* (Guilding, 1828): First record of occurrence for the Northwestern region of Argentina. *Check List* 7(3): 263-266.
- Richardot, M. 1974. La forme septifere de *Ferrissia wautieri* (Moll. Basomm. Ancyliidae). *Haliotis* 2: 203-204.
- Santos, S.B. 2003. Estado atual do conhecimento dos ancilídeos na América do Sul. (Mollusca, Gastropoda, Pulmonata, Basommatophora). *Revista de Biologia Tropical* 50(Suppl. 3): 191-224.
- Simone, R.L., T.C. Bunioto, W.E.P. Avelar and C. Hayashi. 2012. Morphology and biological aspects of *Gundlachia ticaga* from S.E. Brazil (Gastropoda: Basommatophora: Ancyliidae) *Archiv für Molluskenkunde* 141(1): 21-30.
- Thiengo, S.C., A.C. Mattos, M.F. Boaventura and M.A. Fernandez. 2004a. Freshwater snails and schistosomiasis mansoni in the state of Rio de Janeiro, Brazil: IV - Sul Fluminense mesoregion. *Memórias do Instituto Oswaldo Cruz* 99: 275-280.
- Thiengo S.C., A.C. Mattos, M.F. Boaventura, M.S. Loureiro, S.B. Santos and M.A. Fernandez. 2004b. Freshwater snails and schistosomiasis mansoni in the state of Rio de Janeiro, Brazil: V - Norte Fluminense Mesoregion. *Memórias do Instituto Oswaldo Cruz* 99(Suppl.1): 99-103.
- Thiengo, S.C., A.C. Mattos, S.B. Santos and M.A. Fernandez. 2006. Freshwater snails and schistosomiasis mansoni in the state of Rio de Janeiro, Brazil: VI - Noroeste Fluminense Mesoregion. *Memórias do Instituto Oswaldo Cruz* 101: 239-245.
- Thiengo, S.C., M.A. Fernandez, M.F. Boaventura, S.B. Santos and A.C. Mattos. 2002a. Freshwater snails and schistosomiasis mansoni in the state of Rio de Janeiro, Brazil: II - Centro Fluminense Mesoregion. *Memórias do Instituto Oswaldo Cruz* 97: 621-626.
- Thiengo, S.C., M.A. Fernandez, M.F. Boaventura, M.G. Magalhães and S.B. Santos. 2002b. Freshwater snails and schistosomiasis mansoni in the state of Rio de Janeiro, Brazil: III - Baixadas Mesoregion. *Memórias do Instituto Oswaldo Cruz* 97(Suppl. I): 43-46.
- Thiengo S.C., M.A. Fernandez, M.F. Boaventura, C.E. Grault, H.F.R. Silva, A.C. Mattos and S.B. Santos. 2001. Freshwater snails and schistosomiasis mansoni in the state of Rio de Janeiro, Brazil: I - Metropolitan Mesoregion. *Memórias do Instituto Oswaldo Cruz* 96(Suppl.): 177-184.
- Thiengo, S.C., S.B. Santos and M.A. Fernandez. 2005. Malacofauna límnic da área de influência do lago da usina hidrelétrica de Serra da Mesa, Goiás, Brasil. I. Estudo qualitativo. *Revista Brasileira de Zoologia* 22(4): 867-874.

RECEIVED: August 2011

ACCEPTED: November 2012

PUBLISHED ONLINE: March 2013

EDITORIAL RESPONSIBILITY: Robert Forsyth